

Plane Symmetric Solutions of Gravitational Field Equations in Five Dimensions

A. N. Aliev*

Feza Gürsey Institute, 34684 Çengelköy, İstanbul, Turkey

H. Cebeci†

Department of Physics, Anadolu University, 26470 Eskişehir, Turkey

T. Dereli‡

Department of Physics, Koç University, 34450 Sarıyer-İstanbul, Turkey

Abstract

We present the effective field equations obtained from a generalized gravity action with Euler-Poincaré term and a cosmological constant in a D dimensional bulk space-time. A class of plane-symmetric solutions that describe a 3-brane world embedded in a $D = 5$ dimensional bulk space-time are given.

Paper presented at the **5th Workshop on Quantization, Dualities and Integrable Systems**, Pamukkale University, 23-28 January 2006, Denizli, Turkey.

*E.mail: aliev@gursey.gov.tr

†E.mail: cebeci@gursey.gov.tr

‡E.mail: tdereli@ku.edu.tr

1 Introduction

Brane-world theories that receive a lot of interest recently are strictly motivated by string models [1]. They were mainly proposed to provide new solutions to the hierarchy problem and compactification of extra dimensions [2],[3]. The main content of the brane-world idea is that we live in a four dimensional world embedded in a higher dimensional bulk space-time. According to the brane-world scenarios, the gauge fields, fermions and scalar fields of the Standard Model should be localised on a 3-brane, while gravity may freely propagate into the higher dimensional bulk.

In our previous work [4] we derived covariant gravitational field equations on a 3-brane embedded in a five-dimensional bulk space-time with $\mathbb{Z}_2$ symmetry in a generalization that included a dilaton scalar as well as the second order Euler-Poincaré density in the action. We introduced a general ADM-type coordinate setting to show that the effective gravitational field equations on the 3-brane remain unchanged, however, the evolution equations off the brane are significantly modified due to the acceleration of normals to the brane surface in the non-geodesic, ADM slicing of space-time.

In the second part of this paper, using the language of differential forms, we present the field equations of a generalized gravity model with a dilaton 0-form and an axion 3-form in Einstein frame from an action that includes the second order Euler-Poincaré term and a cosmological constant in a D -dimensional bulk space-time. In the third part, we present some plane-symmetric solutions that generalize the well-known domain-wall solution [5].

2 Model

We consider a D -dimensional bulk space-time manifold M equipped with a metric g and a torsion-free, metric compatible connection ∇ . We determine our gravitational field equations by a variational principle from a D -dimensional action that includes the second order Euler-Poincaré term and a cosmological constant

$$I[e, \omega, \phi, H] = \int_M \mathcal{L} \tag{2.1}$$

where in the Einstein frame the Lagrangian density D-form [6]

$$\begin{aligned}\mathcal{L} = & \frac{1}{2}R^{ab} \wedge *(e_a \wedge e_b) - \frac{\alpha}{2}d\phi \wedge *d\phi + \frac{\beta}{2}e^{-\beta_2\phi}H \wedge *H + \Lambda e^{-\beta_1\phi} * 1 \\ & + \frac{\eta}{4}R^{ab} \wedge R^{cd} \wedge *(e_a \wedge e_b \wedge e_c \wedge e_d) \\ & + (de^a + \omega^a{}_b \wedge e^b) \wedge \lambda_a + (dH - \frac{\varepsilon}{2}R_{ab} \wedge R^{ab}) \wedge \mu .\end{aligned}\quad (2.2)$$

Here λ_a and μ are Lagrange multiplier forms that upon variation impose the zero-torsion and anomaly-freedom constraints.

The final form of the variational field equations to be solved are the Einstein field equations

$$\begin{aligned}\frac{1}{2}R^{ab} \wedge *(e_a \wedge e_b \wedge e_c) = & -\frac{\alpha}{2}\tau_c[\phi] + \frac{\beta}{2}e^{-\beta_2\phi}\tau_c[H] - \Lambda e^{-\beta_1\phi} * e_c \\ & - \frac{\eta}{4}R^{ab} \wedge R^{dg} \wedge *(e_a \wedge e_b \wedge e_d \wedge e_g \wedge e_c) \\ & - 2\varepsilon\beta D(e^{-\beta_2\phi}\iota_b(R^b{}_c \wedge *H)) - \frac{\varepsilon\beta}{2}e_c \wedge D(e^{-\beta_2\phi}\iota_s\iota_l(R^{ls} \wedge *H)),\end{aligned}\quad (2.3)$$

where the dilaton stress-energy forms

$$\tau_a[\phi] = \iota_a d\phi * d\phi + d\phi \wedge \iota_a * d\phi$$

and the axion stress-energy forms

$$\tau_a[H] = \iota_a H \wedge *H + H \wedge \iota_a * H ,$$

the dilaton scalar field equation

$$\alpha d(*d\phi) = \frac{\beta_2\beta}{2}e^{-\beta_2\phi}H \wedge *H + \Lambda\beta_1 e^{-\beta_1\phi} * 1, \quad (2.4)$$

and the axion field equations

$$dH = \frac{\varepsilon}{2}R_{ab} \wedge R^{ab} \quad , \quad d(e^{-\beta_2\phi} * H) = 0. \quad (2.5)$$

3 Plane symmetric solutions in $D = 5$

We investigate below a class of plane symmetric solutions in 5-dimensions. We consider the metric

$$g = -f^2(t, \omega)dt^2 + u^2(t, \omega)d\omega^2 + g^2(t, \omega) \left(\frac{dx^2 + dy^2 + dz^2}{(1 + \frac{kr^2}{4})^2} \right), \quad (3.1)$$

the dilaton scalar field

$$\phi = \phi(t, \omega) \quad (3.2)$$

and 3-form gauge field

$$H = h(t, \omega) \frac{dx \wedge dy \wedge dz}{\left(1 + \frac{kr^2}{4}\right)^3} \quad (3.3)$$

in terms of local coordinates

$$x^M : \{x^0 = t, x^5 = \omega, x^1 = x, x^2 = y, x^3 = z\}.$$

We choose our co-frame 1-forms as

$$e^0 = f(t, \omega)dt, \quad e^5 = u(t, \omega)d\omega, \quad e^i = g(t, \omega) \frac{dx^i}{\left(1 + \frac{kr^2}{4}\right)}, \quad i = 1, 2, 3. \quad (3.4)$$

Then we calculate the Levi-Civita connection 1-forms

$$\omega^0{}_i = \frac{g_t}{fg}e^i, \quad \omega^i{}_j = \frac{k}{2g}(x^i e^j - x^j e^i), \quad (3.5)$$

$$\omega^0{}_5 = \frac{u_t}{fu}e^5 + \frac{f_\omega}{fu}e^0, \quad \omega^i{}_5 = \frac{g_\omega}{ug}e^i. \quad (3.6)$$

and the corresponding curvature 2-forms

$$R^{ij} = \frac{1}{g^2} \left\{ k + \left(\frac{g_t}{f}\right)^2 - \left(\frac{g_\omega}{u}\right)^2 \right\} e^i \wedge e^j, \quad (3.7)$$

$$R^{05} = \frac{1}{fu} \left\{ \left(\frac{f_\omega}{u}\right)_\omega - \left(\frac{u_t}{f}\right)_t \right\} e^5 \wedge e^0, \quad (3.8)$$

$$R^{0i} = \frac{1}{fg} \left\{ \left(\frac{g_t}{f}\right)_t - \frac{f_\omega g_\omega}{u^2} \right\} e^0 \wedge e^i + \frac{1}{ug} \left\{ \left(\frac{g_t}{f}\right)_\omega - \frac{u_t g_\omega}{fu} \right\} e^5 \wedge e^i, \quad (3.9)$$

$$R^{i5} = \frac{1}{fg} \left\{ \frac{f_\omega g_t}{fu} - \left(\frac{g_\omega}{u}\right)_t \right\} e^i \wedge e^0 + \frac{1}{ug} \left\{ \left(\frac{g_\omega}{u}\right)_\omega - \frac{g_t u_t}{f^2} \right\} e^5 \wedge e^i. \quad (3.10)$$

From these expressions we note that $R_{ab} \wedge R^{ab} = 0$. Therefore $dH = 0$ implying that

$$H = \frac{Q}{g^3} e^1 \wedge e^2 \wedge e^3 \quad (3.11)$$

where Q may be identified as a magnetic charge. Now, for simplicity, we let $k = 0$ and take the functions g , f and u independent of time. Then we obtain the following system of coupled ordinary differential equations ($'$ denotes derivative with respect to ω):

$$2G - 2C - B - A = -\eta(2CG - AB) - \frac{\alpha}{2} \left(\frac{\phi'}{u} \right)^2 - \frac{\beta Q^2}{2 g^6} e^{-\beta_1 \phi} + \Lambda e^{-\beta_2 \phi}, \quad (3.12)$$

$$3A - 3G = 3\eta GA + \frac{\alpha}{2} \left(\frac{\phi'}{u} \right)^2 - \frac{\beta Q^2}{2 g^6} e^{-\beta_2 \phi} - \Lambda e^{-\beta_1 \phi}, \quad (3.13)$$

$$3C + 3A = -3\eta CA - \frac{\alpha}{2} \left(\frac{\phi'}{u} \right)^2 - \frac{\beta Q^2}{2 g^6} e^{-\beta_2 \phi} - \Lambda e^{-\beta_1 \phi}, \quad (3.14)$$

$$\alpha \left(\frac{\phi' f g^3}{u} \right)' \frac{1}{g^3 f u} = \frac{\beta_2 \beta}{2} e^{-\beta_2 \phi} \frac{Q^2}{g^6} + \Lambda \beta_1 e^{-\beta_1 \phi}. \quad (3.15)$$

where

$$A = - \left(\frac{g'}{g} \right)^2 \frac{1}{u^2} \quad B = - \left(\frac{f'}{u} \right)' \frac{1}{f u}, \quad (3.16)$$

$$C = - \frac{f' g'}{u^2 f g} \quad G = \left(\frac{g'}{u} \right)' \frac{1}{u g}. \quad (3.17)$$

We will give below some special classes of solutions:

Case: $\phi = \text{constant}$, $H = 0$ and $\eta = 0$.

Here the Euler-Poincaré term is absent, $H = 0$ and the dilaton scalar is constant. We obtain the AdS solution in 5-dimensions that is also known as Randall-Sundrum model [3]:

$$g = d\omega^2 + e^{\mp 2p\omega} (-dt^2 + dx^2 + dy^2 + dz^2). \quad (3.18)$$

where $p^2 = \frac{\Lambda}{6}$.

Case: $\phi = \text{constant}$, $H = 0$.

Here $H = 0$ and the dilaton scalar is constant. Solutions are given by the metric

$$g = d\omega^2 + e^{\mp 2s\omega} (-dt^2 + dx^2 + dy^2 + dz^2) \quad (3.19)$$

where

$$s^2 = \frac{1 + \sqrt{1 - \frac{\eta\Lambda}{3}}}{\eta} \quad (3.20)$$

provided that $\Lambda\eta \leq 3$. When $\eta\Lambda = 3$, the solution may alternatively be given in AdS form as

$$g = -4 \cosh^2(l\omega) dt^2 + d\omega^2 + 4 \sinh^2(l\omega) (dx^2 + dy^2 + dz^2) \quad (3.21)$$

where $l^2 = \frac{1}{\eta}$.

Case: $\eta = 0$, $H = 0$.

Here the Euler-Poincaré term is absent and $H = 0$. We obtain the following solution:

$$g = e^{\frac{16\alpha}{3\beta_1}\phi(\omega)} d\omega^2 + e^{\frac{4\alpha}{3\beta_1}\phi(\omega)} (-dt^2 + dx^2 + dy^2 + dz^2) \quad (3.22)$$

with

$$\phi(\omega) = \frac{1}{\left(\frac{\beta_1}{2} - \frac{8\alpha}{3\beta_1}\right)} \ln \left| \sqrt{\frac{2\beta_1\Lambda}{\left(\frac{16\alpha}{3\beta_1} - \beta_1\right)\alpha}} \left(\frac{\beta_1}{2} - \frac{8\alpha}{3\beta_1}\right) \omega + C_0 \right| \quad (3.23)$$

where C_0 is an integration constant. When $\beta_1 = 2$, it reduces to a supersymmetric domain wall solution presented in [5].

Case: $\eta = 0$.

In this case the solution possesses a magnetic charge. It is given by

$$g = e^{\frac{4(\beta_1 - \beta_2)}{3}\phi(\omega)} d\omega^2 + e^{\frac{(\beta_1 - \beta_2)}{3}\phi(\omega)} (-dt^2 + dx^2 + dy^2 + dz^2) \quad (3.24)$$

with

$$\phi(\omega) = \frac{6}{4\beta_2 - \beta_1} \ln \left| \left(\frac{4\beta_2 - \beta_1}{6}\right) \sqrt{\frac{6\left(\frac{\beta_2\beta}{2}Q^2 + \Lambda\beta_1\right)}{(\beta_1 - 4\beta_2)\alpha}} \omega + C \right| \quad (3.25)$$

provided that the constants satisfy

$$(\beta_1 - \beta_2) \left(\frac{\beta Q^2 \beta_2}{2} + \beta_1 \Lambda \right) = \left(\frac{\beta Q^2}{2} + 4\Lambda \right) \alpha. \quad (3.26)$$

C is an integration constant. H is given by

$$H = Qe^{\frac{(\beta_2 - \beta_1)}{2}\phi(\omega)} e^1 \wedge e^2 \wedge e^3 \quad (3.27)$$

We note that when $Q = 0$ and the constants β_1 and β_2 satisfy $\beta_1 - \beta_2 = \frac{4\alpha}{\beta_1}$, the solutions reduce to (3.22) and (3.23).

We also note that an electric dual of solutions (3.24) and (3.25) may be given by defining a 2-form field

$$F = e^{\beta_2\phi} * H. \quad (3.28)$$

Then the solutions are identified as electrically charged solutions.

4 Conclusion

We have given a class of solutions to the variational field equations of a generalized theory of gravity in a D dimensional bulk space-time derived from an action that includes the second-order Euler-Poincaré term and a cosmological constant. The theory describes a heterotic type first order effective string model in D dimensions in the Einstein frame. The special class of plane-symmetric solutions of this model in 5-dimensions we gave refer to a 3-brane world also called a domain wall solution in the literature [5].

References

- [1] P. Horava and E. Witten, Nucl. Phys. B **475**, (1996), 94.
- [2] N. Arkani-Hamed, S. Dimopoulos and G. Dvali, Phys. Lett. B **429**, (1998), 263,
I. Antoniadis, N. Arkani-Hamed , S. Dimopoulos and G. Dvali, Phys. Lett. B **436**, (1998), 257.
- [3] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, (1999), 3370,
L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, (1999), 4690.
- [4] A. N. Aliev, H. Cebeci and T. Dereli, Class. Quant. Grav. **23**, (2006), 591.
- [5] A. Lukas, B. A. Ovrut, K. S. Stelle and D. Waldram, Phys. Rev. D **59**, (1999), 086001.
- [6] T. Dereli and R. W. Tucker, Class. Quant. Grav. **4**, (1987), 791.